



## OPEN ACCESS

## EDITED BY

Boris Klempa,  
Slovak Academy of Sciences, Slovakia

## REVIEWED BY

Vanessa Herder,  
MRC-University of Glasgow Centre For  
Virus Research (MRC), United Kingdom  
Luboš Korytár,  
University of Veterinary Medicine and  
Pharmacy in Košice, Slovakia

## \*CORRESPONDENCE

Joseph A. Jackson,  
✉ j.a.jackson@salford.ac.uk

RECEIVED 24 April 2024

ACCEPTED 26 July 2024

PUBLISHED 08 August 2024

## CITATION

Jackson JA (2024), A novel morbillivirus  
and a novel betaherpesvirus infecting  
the Wood Mouse in the UK.  
*Acta Virol.* 68:13177.  
doi: 10.3389/av.2024.13177

## COPYRIGHT

© 2024 Jackson. This is an open-access  
article distributed under the terms of the  
[Creative Commons Attribution License  
\(CC BY\)](#). The use, distribution or  
reproduction in other forums is  
permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original  
publication in this journal is cited, in  
accordance with accepted academic  
practice. No use, distribution or  
reproduction is permitted which does  
not comply with these terms.

# A novel morbillivirus and a novel betaherpesvirus infecting the Wood Mouse in the UK

Joseph A. Jackson\*

School of Science, Engineering and Environment, University of Salford, Salford, United Kingdom

A novel morbillivirus and a novel betaherpesvirus are reported in the Wood Mouse (*Apodemus sylvaticus*) in the western United Kingdom (UK). The two viruses were found coinfecting an underweight host with abnormalities of the liver and were detected via deep sequencing of lung RNA and *de novo* assembly of substantial genome fragments. The phylogenetic affinities of the novel viruses are characterised based on their relationships to existing database sequences.

## KEYWORDS


*Apodemus sylvaticus*, betaherpesvirus, liver disease, morbillivirus, pulmonary

## Introduction

The virome of wild rodents in the UK, and globally, is of intrinsic ecological importance and also of great significance as a reservoir from which zoonotic and trans-species infectious diseases may emerge (Wu et al., 2018). The viral burden of UK wildlife is only partially documented but a more complete picture would be of value in assessing public and animal health risks (Carlson et al., 2021) and in understanding the regulation and health of wild populations (Lochmiller, 1996). In this report two novel viruses are documented from a wild rodent, the Wood Mouse (*Apodemus sylvaticus*), in the UK. These viruses were discovered via *de novo* assembly of substantial genomic fragments, working from Illumina short-read sequences of host lung RNA. The context of the discovery indicates the likelihood of substantial undiscovered diversity.

## Method

A single mature male *A. sylvaticus* was captured in a forestry plantation near the confluence of the Afon Tarennig and the Afon Gwy/River Wye (52.426773, -3.707703) in July 2011 by humane live-trapping and killed via a UK Home Office approved method. In the UK, *Apodemus* mice are represented only by *A. sylvaticus* and *A. flavicollis* (see Mathews et al., 2018). The latter is not known to occur in the study area, although it can occur in adjacent areas of Mid Wales (Mathews et al., 2018). To avoid any possibility of misidentification, the present specimen was initially positively identified on the basis of morphological criteria, primarily ventral colouration and foot length (Flowerdew, 1984). This identification was consistent with *de novo* assembled host transcripts derived from



**FIGURE 1**


Neighbour joining tree (Saitou and Nei, 1987) of representative morbillivirus nucleocapsid proteins, based on 388 amino acid positions aligned using ClustalW (Thompson et al., 1994). Percentage bootstrap values supporting clusters are shown next to corresponding branches. GenBank accession numbers and a taxon name are shown for each sequence. The scale bar shows amino acid substitutions per site. Phylogenetic analysis was conducted in MEGA Version 10.0.05 (Kumar et al., 2018).

the RNAseq data that are described further below. For example, 840 base pair fragments of coding sequences for *A. flavicollis* toll-like receptor 5 (*tlr5*) haplotypes in National Center for Biotechnology Information (NCBI) GenBank (Sayers et al., 2022) (GenBank accession numbers: OM365774-6) share 98.2%–98.3% identity with a corresponding transcript fragment for the present specimen. In comparison, this same fragment from the present specimen shares 99.5% identity with a homologous fragment predicted from the *A. sylvaticus* NCBI (Sayers et al., 2022) RefSeq genome (XM\_052201127). The captured specimen was underweight, with a negative residual of c.2.5 g from a weight on length regression of mature males ( $n = 12$ ) measured at nearby sites in summer 2011. Furthermore, upon internal examination, there were multiple adhesions of the liver to the abdominal cavity wall and abnormal, fibrotic growth of liver tissue investing the anteroventral surface of the right kidney. No other abnormalities were observed. Fresh, apparently healthy, lung tissue was removed and stored in RNA stabilization solution. Following total RNA extraction from this material and poly (A)-focussed sequencing library construction (originally intended as a control for a separate study, and not aimed primarily at virus discovery in the present specimen), the library was sequenced on an Illumina Novaseq machine (at Azenta Life Sciences, UK) yielding c. 171 million 150 base pair paired end reads. The reads were mapped to a *Mus musculus* genome (mm9) using *BBMap* (Bushnell et al., 2017)

and unmapped reads were *de novo* assembled employing *rnairalSPAdes* (Meleshko and Korobeynikov, 2023). Assembled contigs were then searched against the NCBI (Sayers et al., 2022) RefSeq viral genomes database via discontinuous megablast (Camacho et al., 2009) at a 40% identity cut-off, with only the strongest hit per contig considered. Substantial alignments (>200 base pairs) to viral sequences were further investigated via individual *blastn* (Camacho et al., 2009) searches against the NCBI (Sayers et al., 2022) nt database and only unambiguous matches to viral sequences (excluding phages and retroviruses) were considered further.

## Results and discussion

We found clear evidence for the presence of two novel viruses in the lungs of the Wood Mouse from our study site. In one case, two large genome fragments were assembled of 1684 (GenBank accession number: PP188092) and 1592 base pairs (PP188093). These unambiguously clustered with rodent-infecting members of the *Morbillivirus* genus in phylogenetic analyses (see Figure 1). Classical morbilliviruses occur in larger mammals and are well-known agents of serious diseases, such as measles, canine distemper, phocine distemper, rinderpest, peste des petits ruminants and cetacean morbillivirus (Seki and Takeda, 2022;



**FIGURE 2**

Neighbour joining tree (Saitou and Nei, 1987) of representative betaherpesvirus m142 and related proteins, based on 194 amino acid positions aligned using ClustalW (Thompson et al., 1994). Percentage bootstrap values supporting clusters are shown next to corresponding branches. GenBank accession numbers and a protein identifier/taxon name are shown for each sequence. The scale bar shows amino acid substitutions per site. Phylogenetic analysis was conducted in MEGA Version 10.0.05 (Kumar et al., 2018).

Libbey and Fujinami, 2023). The present discovery adds to the divergent members of the *Morbillivirus* genus that have recently been described in rodents (Debat, 2022; Vanmechelen et al., 2022; Chen et al., 2023) and bats (Wells et al., 2022; Ikegame et al., 2023). The novel morbillivirus (Tarennig *A. sylvaticus* morbillivirus) is closest to, and might be considered a well differentiated strain of, Gierle Apodemus virus (Vanmechelen et al., 2022). The 1684 base pair assembled fragment encompassed a full 1572 base pair open reading frame coding for a 524 amino acid nucleocapsid-like protein with 95% residue similarity to the 526 amino acid nucleocapsid protein (UQM99541.1) in Gierle Apodemus Virus. The 1592 base pair fragment encompassed a full 1488 base pair open reading frame coding for a 496 amino acid phosphoprotein-like protein with 89% residue similarity to the 496 amino acid phosphoprotein in Gierle Apodemus Virus (UQM99543.1).

In another case, two large genome fragments of 834 base pairs (PP188094) and 546 base pairs (PP188095) were assembled that clustered with murine betaherpesviruses (*Muromegalovirus* (Walker et al., 2022) or *Cytomegalovirus*) in phylogenetic analyses (see Figure 2). The novel virus (Tarennig *A. sylvaticus* betaherpesvirus) was close to, but well differentiated from, Murid betaherpesvirus 1. The larger genome fragment contained an 816 bp part of an open reading frame, corresponding to a 272 amino acid sequence, with greatest

similarity to the m142 protein of Murid betaherpesvirus 1 (AWV68063.1; 67% amino acid identity across a 237 aa aligned region). The smaller genome fragment contained a 543 base pair part of an open reading frame, corresponding to a 181 amino acid sequence, with greatest similarity to the m85 protein of Murid betaherpesvirus 1 and its equivalent (capsid triplex subunit 2) in *Mastomys natalensis* cytomegalovirus 3 (WEG69771.1; with 85% identity over a 158 amino acid aligned region in this case). As far as can be determined this is the first record of a betaherpesvirus in the Wood Mouse in the U.K. or elsewhere, although there is a disputed claim (Kim et al., 1974) that a murine cytomegalovirus derived from Wood Mice was adapted to replicate in human cells *in vitro* in the 1960s (Raynaud et al., 1969). Numerous other betaherpesviruses (Ehlers et al., 2007; Tarlinton et al., 2011; Ntumvi et al., 2018) have been reported in rodents on the basis of small sequence fragments of genes not recovered here, including in *A. flavicollis* (Yellow-necked Mice) in Germany (Ehlers et al., 2007). It was secondarily possible to recover two reads corresponding to the betaherpesvirus DNA polymerase gene, which has been targeted in some viral discovery studies, and these were most similar to *A. flavicollis* cytomegalovirus 2 (EF125063.1; 86%–87% nucleotide identity) and then to Murid betaherpes virus 1 isolates (78%–81% nucleotide identity).

It is recognized that the present study is based on a limited sample (a single tissue from a single host) and that further studies are required to reveal the epidemiological characteristics and biology of the discovered viruses. Nonetheless, the current records usefully extend our knowledge of the virome in UK wild mammals. It is also noted that neither virus was straightforwardly detected via the analysis of short reads with taxonomic classifier softwares under a range of conditions and that such classifiers were prone to false positives due to host-derived rRNA hits. The ready detection of new viruses via the methods employed, even from a single arbitrarily sampled host specimen, and the high level of nucleotide divergence of the viruses from their closest relatives in databases, are suggestive of considerable undiscovered viral diversity.

## Data availability statement

The datasets presented in this study can be found in online repositories. The names of the repository/repository and accession number(s) are as follows: <https://www.ncbi.nlm.nih.gov/genbank/>, PP188092 <https://www.ncbi.nlm.nih.gov/genbank/>, PP188093 <https://www.ncbi.nlm.nih.gov/genbank/>, PP188094 <https://www.ncbi.nlm.nih.gov/genbank/>, PP188095.

## Ethics statement

The study complied with the ethics review procedure at Aberystwyth University at the time the work was carried out

## References

- Bushnell, B., Rood, J., and Singer, E. (2017). BBMerge – accurate paired shotgun read merging via overlap. *Plos one* 12, e0185056. doi:10.1371/journal.pone.0185056
- Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., et al. (2009). BLAST+: architecture and applications. *BMC Bioinform.* 10, 421. doi:10.1186/1471-2105-10-421
- Carlson, C. J., Farrell, M. J., Grange, Z., Han, B. A., Mollentze, N., Phelan, A. L., et al. (2021). The future of zoonotic risk prediction. *Philosophical Trans. R. Soc. Lond. Ser. B, Biol. Sci.* 376, 20200358. doi:10.1098/rstb.2020.0358
- Chen, Y.-M., Hu, S.-J., Lin, X.-D., Tian, J.-H., Lv, J.-X., Wang, M.-R., et al. (2023). Host traits shape virome composition and virus transmission in wild small mammals. *Cell* 186, 4662–4675.e12. doi:10.1016/j.cell.2023.08.029
- Debat, H. J. (2022). A South American mouse morbillivirus provides insight into a clade of rodent-borne morbilliviruses. *Viruses* 14, 2403. doi:10.3390/v14112403
- Ehlers, B., Küchler, J., Yasmum, N., Dural, G., Voigt, S., Schmidt-Chanasit, J., et al. (2007). Identification of novel rodent herpesviruses, including the first gammaherpesvirus of *Mus musculus*. *J. Virol.* 81, 8091–8100. doi:10.1128/jvi.00255-07
- Flowerdew, J. (1984). *Woodmice and yellow-necked mice (mammal society series)*. Oswestry, UK: Nelson.
- Ikegame, S., Carmichael, J. C., Wells, H., Furler O'Brien, R. L., Acklin, J. A., Chiu, H. P., et al. (2023). Metagenomics-enabled reverse-genetics assembly and characterization of myotis bat morbillivirus. *Nat. Microbiol.* 8, 1108–1122. doi:10.1038/s41564-023-01380-4
- Kim, K. S., Sapienza, V., and Carp, R. I. (1974). Comparative studies of the Smith and Raynaud strains of murine cytomegalovirus. *Infect. Immun.* 10, 672–674. doi:10.1128/iai.10.3.672-674.1974
- Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). Mega X: Molecular evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol.* 35, 1547–1549. doi:10.1093/molbev/msy096
- Libbey, J. E., and Fujinami, R. S. (2023). Morbillivirus: A highly adaptable viral genus. *Heliyon* 9, e18095. doi:10.1016/j.heliyon.2023.e18095
- Lochmiller, R. L. (1996). Immunocompetence and animal population regulation. *Oikos* 76, 594–602. doi:10.2307/3546356
- Mathews, F., Kubasiewicz, L. M., Gurnell, J., Harrower, C. A., McDonald, R. A., and Shore, R. F. (2018). *A review of the population and conservation status of British mammals*. Peterborough, UK: Natural England.
- Meleshko, D., and Korobeynikov, A. (2023). Benchmarking state-of-the-art approaches for norovirus genome assembly in metagenome sample. *Biology* 12, 1066. doi:10.3390/biology12081066
- Ntumvi, N. F., Mbala Kingebeni, P., Tamoufe, U., Kumakamba, C., Ndze, V., Ngay Lukusa, I., et al. (2018). High herpesvirus diversity in wild rodent and shrew species in Central Africa. *Intervirology* 61, 155–165. doi:10.1159/000493796
- Raynaud, J., Atanasiu, P., Barreau, C., and Jahkola, M. (1969). Adaptation d'un virus cytomégalique provenant du Mulot (*Apodemus sylvaticus*) sur différentes cellules hétérologues, y compris les cellules humaines. *Comptes Rendus Hebd. Séances L'Académie Sci.* 269, 104–106.

and was conducted in accordance with the relevant local legislation and institutional requirements.

## Author contributions

The author confirms being the sole contributor of this work and has approved it for publication.

## Funding

The author(s) declare that financial support was received for the research, authorship, and/or publication of this article. Fieldwork was supported by Aberystwyth University and sequencing work was supported by the University of Salford.

## Acknowledgments

Grateful acknowledgement is made to the UK Forestry commission, for permission to trap at the field site in 2011, and to the Galaxy Europe server, on which some bioinformatic analyses were carried out.

## Conflict of interest

The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Saitou, N., and Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4, 406–425. doi:10.1093/oxfordjournals.molbev.a040454

Sayers, E. W., Bolton, E. E., Brister, J. R., Canese, K., Chan, J., Comeau, D. C., et al. (2022). Database resources of the national center for biotechnology information. *Nucleic Acids Res.* 50 (D1), D20–D26. doi:10.1093/nar/gkab1112

Seki, F., and Takeda, M. (2022). Novel and classical morbilliviruses: current knowledge of three divergent morbillivirus groups. *Microbiol. Immunol.* 66, 552–563. doi:10.1111/1348-0421.13030

Tarlinton, R. E., Tate, H. F. J., Mohallal, E. M. E., Daly, J. M., and Behnke, J. M. (2011). Betaherpesvirus sequences in eastern spiny mice and wagner's dipodids. *J. Wildl. Dis.* 47, 958–962. doi:10.7589/0090-3558-47.4.958

Thompson, J. D., Higgins, D. G., and Gibson, T. J. (1994). Clustal W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22, 4673–4680. doi:10.1093/nar/22.22.4673

Vanmechelen, B., Meurs, S., Horemans, M., Loosen, A., Joly Maes, T., Laenen, L., et al. (2022). The characterization of multiple novel paramyxoviruses highlights the diverse nature of the subfamily Orthoparamyxovirinae. *Virus Evol.* 8, eac061. doi:10.1093/ve/veac061

Walker, P. J., Siddell, S. G., Lefkowitz, E. J., Mushegian, A. R., Adriaenssens, E. M., Alfenas-Zerbini, P., et al. (2022). Recent changes to virus taxonomy ratified by the international committee on taxonomy of viruses (2022). *Arch. Virol.* 167, 2429–2440. doi:10.1007/s00705-022-05516-5

Wells, H. L., Loh, E., Nava, A., Solorio, M. R., Lee, M. H., Lee, J., et al. (2022). Classification of new morbillivirus and jeilongvirus sequences from bats sampled in Brazil and Malaysia. *Arch. Virol.* 167, 1977–1987. doi:10.1007/s00705-022-05500-z

Wu, Z., Lu, L., Du, J., Yang, L., Ren, X., Liu, B., et al. (2018). Comparative analysis of rodent and small mammal viromes to better understand the wildlife origin of emerging infectious diseases. *Microbiome* 6, 178–214. doi:10.1186/s40168-018-0554-9