

Structure and immunological function of oxidised albumin in lung cancer: its potential role as a biomarker of elevated oxidative stress

Z. RASHEED^{*,‡}, R. AHMAD[†] and R. ALI^{*}

^{*}Department of Biochemistry, Faculty of Medicine, J. N. Medical College, AMU, Aligarh-202002; [†]Department of Biochemistry, SBSPGI, Balawala, Dehradun-248161, India; and [‡]Department of Pathology, Microbiology and Immunology, School of Medicine, University of South Carolina, Columbia, SC-29209, USA

Figure 2 contained an error when this Original Article was published originally (*Br J Biomed Sci* 2009; **66**: 67–73). Figure 2 from this paper is reproduced here in its correct form.




Fig. 2. a) CD spectra of purified HSA from smoking LC patients (▲), non-smoking LC patients (■), healthy smoking subjects (●) and healthy non-smoking subjects (◆). Spectra are the average of eight determinations. b) Fluorescence spectra of purified HSA-bound bis-ANS. Purified HSA from smoking LC patients (▲), non-smoking LC patients (■), healthy smoking subjects (●) and healthy non-smoking subjects (◆). Spectra are the average of eight determinations.

Correspondence to: Dr Zafar Rasheed

Department of Pathology, Microbiology and Immunology, School of Medicine
University of South Carolina, Columbia, SC-29209, USA

Email: zafar.rasheed@uscmcd.sc.edu