

## Clinical and epidemiological characterisation of Burkitt's lymphoma: an eight-year case study at Komfo Anokye Teaching Hospital, Ghana

L. OWUSU\*, F. A YEBOAH<sup>†,‡</sup>, A. OSEI-AKOTO<sup>§,¶</sup>, T. RETTIG<sup>§,¶</sup> and F. K. N. ARTHUR\*

<sup>\*</sup>Departments of Biochemistry and Biotechnology and <sup>†</sup>Molecular Medicine, School of Medical Sciences, Kwame Nkrumah University of Science and Technology; <sup>‡</sup>Departments of <sup>§</sup>Clinical Biochemistry and <sup>¶</sup>Child Health, Komfo Anokye Teaching Hospital; and

<sup>¶</sup>Department of Child Health, School of Medical Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana

Correspondence to:

Fareed K. N. Arthur

Email: fareedarthur@yahoo.com

Figures 4, 5, 10 and 11 were incorrect in the original publication of this article (*Br J Biomed Sci* 2010; **67**: 9–14). The four figures are reproduced here in their correct form and with their correct legends.




Fig. 4. Age and gender distribution of cases.




Fig. 5. Sites of primary tumour presentation by gender.




Fig. 10. Incidence of BL in the regions of Ghana (KATH 2000–2007).




Fig. 11. Regional BL presentation in the regions of Ghana.